

2012 DAGGETT COUNTY NOXIOUS WEED PRIORITY CLASSIFICATION

as reviewed and approved by the Daggett County Weed Board (DCWB) on February 15, 2012. The italicized weeds were moved from Class B to Class A classification.

CLASS A WEEDS

| | |
|---|-------------------------------|
| Black Henbane | <i>Hyoscyamus niger</i> |
| <i>Canada Thistle</i> | <i>Cirsium arvense</i> |
| Dalmation Toadflax | <i>Linaria dalmatica</i> |
| Dyers Woad | <i>Isatis tinctoria</i> L. |
| Leafy Spurge | <i>Euphorbia esula</i> |
| <i>Mountain Thernopsis</i> (<i>mountain pea</i>) | |
| Musk Thistle | <i>Carduus nutans</i> |
| Purple Loosestrife | <i>Lythrum salicaria</i> L. |
| <i>Perennial Pepperweed</i> | <i>Lepidium Latifolium</i> |
| <i>Russian Knapweed</i> | <i>Centaurea repens</i> |
| Scotch Thistle | <i>Onopordum Acanthium</i> |
| Yellow Starthistle | <i>Centaurea solstitialis</i> |
| Yellow Toadflax | <i>Linaria vulgaris</i> |

CLASS B WEEDS

| | |
|-------------------------------|-------------------------------|
| Common Burdock | |
| Diffused Knapweed | <i>Centaurea diffusa</i> |
| Hoary cress (short white top) | <i>Cardaria</i> Spp |
| Houndstongue | <i>Cynaglossum officinale</i> |
| Saltcedar | <i>Tamarix ramosissima</i> |
| Spotted Knapweed | <i>Centaurea maculosa</i> |

CLASS C WEEDS

| | |
|--|-----------------------------|
| Field Bindweed (Wild Morning Glory) | <i>Convolvulus arvensis</i> |
|--|-----------------------------|

“Class A” weeds have a relatively low population size within the State and are the highest priority being an *Early Detection Rapid Response* (EDRR) weed.

“Class B” weeds have a moderate population throughout the State and generally are thought to be controllable in most areas.

“Class C” weeds are found extensively in the State and are thought to be beyond control. Statewide efforts would generally be towards containment of smaller infestations.